


Monthly Geomagnetic Bulletin

February 2003 03/02/HA



1. HARTLAND OBSERVATORY MAGNETIC DATA

1.1 Introduction

This bulletin is published to meet the needs of both commercial and academic users of geomagnetic data. Magnetic observatory data is presented as a series of plots of one-minute, hourly and daily values, followed by tabulations of monthly values, geomagnetic activity indices and reports of rapid variations. The operation of the observatory and presentation of data are described in the rest of this section.

Enquiries about the data should be addressed to:

National Geomagnetic Service
British Geological Survey
Murchison House, West Mains Road
Edinburgh EH9 3LA
Scotland, UK

Tel: +44 (0) 131 667 1000
Fax: +44 (0) 131 668 4368
E-mail: s.reay@bgs.ac.uk
Internet: www.geomag.bgs.ac.uk

1.2 Position

Hartland Observatory, one of the three geomagnetic observatories operated and maintained in the UK by BGS, is situated on the NW boundary of the village of Hartland in North Devon. The observatory co-ordinates are:

Geographic: 50° 59.7' N 355° 31.0' E
Geomagnetic: 53° 55.3' N 80° 8.2' E
Height above mean sea level: 95 m

The geomagnetic co-ordinates are calculated using the 8th generation International Geomagnetic Reference Field at epoch 2003.5.

1.3 The Observatory Operation

1.3.1 GDAS

The observatory operates under the control of the Geomagnetic Data Acquisition System (GDAS), which was developed by BGS staff, installed in 2002, and became fully operational in January 2003. The system operates under the control of data acquisition software running on QNX computers, which control the data-logging and communications.

There are two sets of sensors used for making magnetic measurements. A triaxial linear-core

fluxgate magnetometer, manufactured by the Danish Meteorological Institute, is used to measure the variations in the horizontal (H) and vertical (Z) components of the field. The third sensor is oriented perpendicular to these, and measures variations, which are proportional to the changes in declination (D). Measurements are made at a rate of 1 Hz.

In addition to the fluxgate sensors there is a proton precession magnetometer making measurements of the absolute total field intensity (F) at a rate of 0.1 Hz.

The raw unfiltered data are retrieved automatically via Internet connections to the BGS office in Edinburgh in near real-time. The fluxgate data are filtered to produce one-minute values using a 61-point cosine filter whilst the total field intensity samples are filtered using a 7-point cosine filter. These one-minute values are used to update the Geomagnetism Information and Forecast Service (GIFS), an on-line information system accessed via the World Wide Web at the address given in Section 1.1. GIFS also provides information on geomagnetic and solar activity.

1.3.2 Back-up Systems

There are two other fully independent identical systems, GDAS 2 and GDAS 3, operating at the observatory. The data from these are also processed in near real-time and used for quality control purposes. They can be also be used to fill any gaps or replace any corrupt values in the primary system, GDAS 1.

1.4 Data Presentation

The data presented in the bulletin are in the form of plots and tabulations described in the following sections.

1.4.1 Summary magnetograms

Small-scale magnetograms are plotted which allow the month's data to be viewed at a glance. They are plotted 16 days a page and show the variations in D , H and Z . The scales are shown on the right-hand side of the page. On disturbed days the scales are multiplied by a factor, which is indicated above the panel for that day. The variations are centred on the monthly mean value, shown on the left side of the page.

1.4.2 Magnetograms

The daily magnetograms are plotted using one-minute values of D , H and Z from the fluxgate sensors, with any gaps filled using back-up data. The magnetograms are plotted to a variable scale; scale bars are shown to the right of each plot. The absolute level (the monthly mean value) is indicated on the left side of the plots.

1.4.3 Hourly Mean Value Plots

Hourly mean values of D , H and Z for the past 12 months are plotted in 27-day segments corresponding to the Bartels solar rotation number. Magnetic disturbances associated with active regions on the surface of the Sun may recur after 27 days: the same is true for geomagnetically quiet intervals. Plotting the data in this way highlights this recurrence, and also illustrates seasonal and diurnal variations throughout the year.

1.4.4 Daily and Monthly Mean Values

Daily mean values of D , H , Z and F are plotted throughout the year. In addition, a table of monthly mean values of all the geomagnetic elements is provided. These values depend on accurate specification of the fluxgate sensor baselines. Provisional and definitive values are indicated in the table as **P** or **D** respectively. It is anticipated that provisional values will not be altered by more than a few nT or tenths of arcminutes before being made definitive.

1.4.5 Geomagnetic activity indices

The Observatory K index. This summarises geomagnetic activity at an observatory by assigning a code, an integer in the range 0 to 9, to each 3-hour Universal Time (UT) interval. The index for each 3-hour UT interval is determined from the ranges in H and in D (scaled in nT), with allowance made for the regular (undisturbed) diurnal variation. The conversion from range to an index value is made using a quasi-logarithmic scale, with the scale values dependent on the geomagnetic latitude of the observatory. The K index retains the local time (LT) and seasonal dependence of activity associated with the position of the observatory.

The provisional aa index. A number of 3-hour geomagnetic indices are computed by combining K indices from networks of observatories to characterise global activity levels and to eliminate LT and seasonal effects. The simplest of these is the aa index, computed using the K indices from two approximately antipodal observatories: Hartland in the UK and Canberra in Australia. The aa index is calculated from linearisations of the

Hartland and Canberra K indices, and has units of nT. The daily mean value of aa (denoted Aa), the mean values of aa for the intervals 00-12UT and 12-24UT and the daily mean values for Hartland alone (Aa_n) and Canberra alone (Aa_s) are tabulated.

Although the aa index is based on data from only two observatories, provided averages over 12 hours or longer are used, the index is strongly correlated with the ap and am indices, which are derived using data from more extensive observatory networks.


The aa indices listed in this publication are provisional only; the definitive values are published by the International Service for Geomagnetic Indices, CRPE/CNET - CNRS, 4 Avenue de Neptune, F-94107 Saint Maur Cedex, France.


1.4.6 Rapid Variations

Charged particles stream from the Sun in the solar wind. The solar wind interacts with the geomagnetic field to create a cavity, the magnetosphere, in which the field is confined. When a region of enhanced velocity and/or density in the solar wind arrives at the day-side boundary of the magnetosphere (at about 10 earth radii) the boundary is pushed towards the Earth. Currents set up on the boundary of the magnetosphere can cause an abrupt change in the geomagnetic field measured on the ground and this is recorded on observatory magnetograms as a Sudden Impulse (SI). If, following an SI, there is a change in the rhythm of activity, the SI is termed a Storm Sudden Commencement (SSC). A classical magnetic storm exhibiting initial, main and recovery phases (shown by, for instance, the Dst ring current index) can often occur after a SSC, in which case the start of the storm is taken as the time of the SSC.

Solar flares, seen at optical wavelengths as a sudden brightening of a small region of the Sun's surface, are also responsible for increased X-ray emissions. The X-rays cause increased ionisation in the ionosphere, which leads to absorption of short-wave radio signals. On an observatory magnetogram a Solar Flare Effect (SFE), or "crochet" may be observed. This is an enhancement to the diurnal variation of the order of 10 nT, lasting about an hour.

Hartland February 2003





Hartland


February

2003

Date: 01-02-2003


Hartland

Day number: 032



Date: 02-02-2003


Day number: 033



Date: 03-02-2003


Hartland

Day number: 034



Date: 04-02-2003


Day number: 035



Date: 05-02-2003


Hartland

Day number: 036



Date: 06-02-2003


Day number: 037



Date: 07-02-2003


Hartland

Day number: 038



Date: 08-02-2003


Day number: 039



Date: 09-02-2003


Hartland

Day number: 040



Date: 10-02-2003


Day number: 041



Date: 11-02-2003


Hartland

Day number: 042



Date: 12-02-2003


Day number: 043



Date: 13-02-2003


Hartland

Day number: 044



Date: 14-02-2003


Day number: 045



Date: 15-02-2003


Hartland

Day number: 046



Date: 16-02-2003


Day number: 047



Date: 17-02-2003


Hartland

Day number: 048



Date: 18-02-2003


Day number: 049



Date: 19-02-2003


Hartland

Day number: 050



Date: 20-02-2003


Day number: 051



Date: 21-02-2003


Hartland

Day number: 052



Date: 22-02-2003


Day number: 053



Date: 23-02-2003


Hartland

Day number: 054



Date: 24-02-2003


Day number: 055



Date: 25-02-2003


Hartland

Day number: 056



Date: 26-02-2003


Day number: 057



Date: 27-02-2003


Hartland

Day number: 058




Date: 28-02-2003


Day number: 059




Hartland Observatory: Declination (degrees)




Hartland Observatory: Horizontal Intensity (nT)



Hartland Observatory: Vertical Intensity (nT)



Hartland 2003



Monthly Mean Values for Hartland Observatory 2003

Month	<i>D</i>	<i>H</i>	<i>I</i>	<i>X</i>	<i>Y</i>	<i>Z</i>	<i>F</i>	Data
January	-4° 19.2'	19542 nT	66° 6.4'	19486 nT	-1472 nT	44114 nT	48249 nT	P
February	-4° 18.1'	19540 nT	66° 6.7'	19485 nT	-1466 nT	44118 nT	48252 nT	P

INDICES OF GEOMAGNETIC ACTIVITY

The K Index

Hartland Observatory

February 2003

Day	K - INDICES FOR THREE-HOUR INTERVAL								SUM
	00-03	03-06	06-09	09-12	12-15	15-18	18-21	21-24	
1	3	2	0	0	1	2	5	5	18
2	5	4	4	4	4	5	5	5	36
3	4	3	2	2	3	4	4	4	26
4	4	4	5	4	2	3	4	4	30
5	3	2	2	3	3	4	2	3	22
6	2	1	2	3	4	3	4	3	22
7	3	3	3	2	3	2	4	3	23
8	3	3	2	3	3	2	3	4	23
9	3	3	3	2	4	4	3	3	25
10	3	4	3	1	2	1	3	4	21
11	4	2	2	2	1	1	1	1	14
12	1	2	1	2	2	4	4	4	20
13	2	1	1	2	1	1	2	4	14
14	3	3	2	3	3	3	4	4	25
15	4	4	3	4	3	4	2	3	27
16	3	3	2	2	3	3	4	4	24
17	3	1	3	3	2	2	2	3	19
18	2	4	3	3	3	2	3	4	24
19	2	2	1	2	3	3	2	4	19
20	2	2	3	4	3	3	4	3	24
21	1	3	3	2	1	4	4	3	21
22	2	3	2	2	3	3	1	3	19
23	3	2	2	2	3	2	3	2	19
24	2	0	1	2	2	1	2	3	13
25	2	1	1	2	2	1	1	1	11
26	1	3	3	3	2	3	3	4	22
27	4	4	3	3	4	4	4	4	30
28	3	1	2	4	3	4	3	4	24

Lower bound (nT) for the range for each index value at Hartland Observatory									
K-Index									
0	1	2	3	4	5	6	7	8	9
0	5	10	20	40	70	120	200	330	500

The aa Index

Date	Day	K-North	K-South	(a)	(b)	(c)	(d)	(e)
01-02-03	32	3 2 0 0 1 2 5 5	1 0 0 0 3 3 4 3	35	22	8	48	28
02-02-03	33	5 4 4 4 4 5 5 5	4 4 6 4 4 5 5 3	79	81	78	82	80
03-02-03	34	4 3 2 2 3 4 4 4	2 2 2 3 4 3 3 3	41	30	25	45	35
04-02-03	35	4 4 5 4 2 3 4 4	3 5 5 4 3 2 3 2	55	49	72	33	52
05-02-03	36	3 2 2 3 3 4 2 3	2 2 2 3 4 3 1 2	29	25	22	32	27
06-02-03	37	2 1 2 3 4 3 4 3	2 2 3 3 3 4 3 3	31	32	21	42	32
07-02-03	38	3 3 3 2 3 2 4 3	2 2 2 2 4 2 3 2	31	24	22	33	27
08-02-03	39	3 3 2 3 3 2 3 4	3 3 3 2 3 3 2 3	31	28	28	31	30
09-02-03	40	3 3 3 2 4 4 3 3	3 2 3 2 3 4 2 3	36	30	26	40	33
10-02-03	41	3 4 3 1 2 1 3 4	2 3 4 3 2 3 2 2	30	28	34	24	29
11-02-03	42	4 2 2 2 1 1 1 1	2 2 2 2 3 3 2 1	17	19	21	15	18
12-02-03	43	1 2 1 2 2 4 4 4	1 2 2 3 2 4 2 2	30	23	15	37	26
13-02-03	44	2 1 1 2 1 1 2 4	1 2 1 2 2 0 1 2	17	11	12	17	14
14-02-03	45	3 3 2 3 3 3 4 4	3 3 2 3 4 4 3 3	36	37	28	46	37
15-02-03	46	4 4 3 4 3 4 2 3	3 4 2 3 4 3 2 3	43	35	43	35	39
16-02-03	47	3 3 2 2 3 3 4 4	2 3 2 3 4 3 2 3	34	30	24	40	32
17-02-03	48	3 1 3 3 2 2 2 3	2 2 2 3 3 2 2 3	23	22	23	22	23
18-02-03	49	2 4 3 3 3 2 3 4	2 6 5 4 3 3 1 3	34	57	61	30	46
19-02-03	50	2 2 1 2 3 3 2 4	2 2 1 2 4 3 2 3	24	25	14	35	24
20-02-03	51	2 2 3 4 3 3 4 3	2 3 4 3 3 2 3 2	34	30	33	31	32
21-02-03	52	1 3 3 2 1 4 4 3	2 2 3 2 1 3 2 2	30	19	21	29	25
22-02-03	53	2 3 2 2 3 3 1 3	1 1 2 2 4 3 2 2	23	22	16	28	22
23-02-03	54	3 2 2 2 3 2 3 2	2 2 1 2 4 3 2 2	22	23	17	27	22
24-02-03	55	2 0 1 2 2 1 2 3	1 1 1 3 2 2 1 2	14	14	12	16	14
25-02-03	56	2 1 1 2 2 1 1 1	1 1 2 1 2 1 0 1	11	9	11	9	10
26-02-03	57	1 3 3 3 2 3 3 4	1 1 3 4 3 2 2 4	30	29	26	33	30
27-02-03	58	4 4 3 3 4 4 4 4	4 3 3 4 5 4 3 3	52	51	46	58	52
28-02-03	59	3 1 2 4 3 4 3 4	3 1 2 4 4 4 3 3	37	38	29	46	37
Monthly mean value =				31.3				

- (a) The northern daily mean value, Aa_n
- (b) The southern daily mean value, Aa_s
- (c) The mean value of aa for the interval 00-12 UT
- (d) The mean value of aa for the interval 12-24 UT
- (e) The daily mean value of aa (Aa)

Notes

- i. The values are rounded to the nearest integer.
- ii. The units of the aa index are nT.
- iii. The values shown here are provisional. The definitive values are computed and published by the International Service for Geomagnetic Indices, Paris

HARTLAND RAPID VARIATIONS

SIs and SSCs

Date	Time (UT)	Type	Quality	H (nT)	D (min)	Z (nT)
01-02-03	13:37	SSC	C	4.4	-0.72	-
18-02-03	04:40	SSC*	B	13.8	3.17	8.2
19-02-03	11:17	SSC	C	4.7	-	1.2

SFEs

Date	Start (UT)	Maximum (UT)	End (UT)	H (nT)	D (min)	Z (Nt)
NONE						

Notes

- i. For SIs and SSCs a * indicates that the principal impulse was preceded by a smaller reversed impulse.
- ii. The quality of SIs and SSCs are classified as follows:
 - A = very distinct
 - B = fair, ordinary, but unmistakable
 - C = doubtful
- iii. The amplitudes given are for the first chief movement of the event.