

BRITISH GEOLOGICAL SURVEY

Port Stanley Observatory Monthly Magnetic Bulletin

March 2012

12/03/PS



**British
Geological Survey**

NATURAL ENVIRONMENT RESEARCH COUNCIL

PORT STANLEY OBSERVATORY MAGNETIC DATA

1. Introduction

Port Stanley observatory was installed by the British Geological Survey (BGS) with financial support from a consortium of oil companies and became operational in February 1994.

This bulletin is published to provide rapid access to the provisional geomagnetic observatory results. The information is freely available for personal, academic, educational and non-commercial research or use. Magnetic observatory data are presented as a series of plots of one-minute, hourly and daily values, followed by tabulations of monthly values. The operation of the observatory and presentation of data are described in the rest of this section.

Enquiries about the data should be addressed to:

Geomagnetism Team
Earth Hazards and Systems
British Geological Survey
Murchison House, West Mains Road
Edinburgh EH9 3LA
Scotland, UK

Tel: +44 (0) 131 667 1000
Fax: +44 (0) 131 650 0265
E-mail: enquiries@bgs.ac.uk
Internet: www.geomag.bgs.ac.uk

2. Position

Port Stanley Observatory, one of the geomagnetic observatories maintained and operated by the British Geological Survey (BGS), is situated on a site at Sapper Hill near Port Stanley in the Falkland Islands. The observatory co-ordinates are:

Geographic: 51° 42' 15" S 302° 06' 24" E
Geomagnetic: 42° 07' 26" S 012° 05' 46" E
Height above mean sea level: 135 m

The geomagnetic co-ordinates are approximations, calculated using the 11th generation International Geomagnetic Reference Field (IGRF) at epoch 2012.5. On-line access to models (including IGRF), charts and navigational data are available at http://www.geomag.bgs.ac.uk/data_service/models_compass/home

3. The Observatory Operation

3.1 GDAS

The observatory operates under the control of the Geomagnetic Data Acquisition System (GDAS), which was developed by BGS staff, installed and became operational in August 2002. The data acquisition software, running on QNX operated computers, controls the data logging and the communications.

There are two sets of sensors used for making magnetic measurements. A tri-axial linear-core fluxgate magnetometer, manufactured by the Danish Meteorological Institute, is used to measure the variations in the horizontal (*H*) and vertical (*Z*) components of the field. The third sensor is oriented perpendicular to these, and measures variations, which are proportional to the changes in declination (*D*). Measurements are made at a rate of 1 Hz.

In addition to the fluxgate sensors there is a proton precession magnetometer (PPM) making measurements of the absolute total field intensity (*F*) at a rate of 0.1Hz.

The raw unfiltered data are retrieved automatically via Internet connections to the BGS office in Edinburgh in near real-time. The fluxgate data are filtered to produce one-minute values using a 61-point cosine filter and the total field intensity samples are filtered using a 7-point cosine filter. The one-minute values provide input for various data products, available on-line at www.geomag.bgs.ac.uk/data_service/home

3.2 Absolute Observations

The GDAS fluxgate magnetometers accurately measure variations in the components of the geomagnetic field, but not the absolute magnitudes. Two sets of absolute measurements of the field are made manually twice per month. A fluxgate sensor mounted on a theodolite is used to determine *D* and inclination (*I*); the GDAS PPM measurements, with a site difference correction applied, are used for *F*. The absolute observations are used in conjunction with the GDAS variometer measurements to produce a continuous record of the absolute values of the geomagnetic field elements as if they had been measured at the observatory reference pillar.

4. Observatory Results

The data presented in the bulletin are in the form of plots and tabulations described in the following sections.

4.1 Absolute Observations

The absolute observation measurements made during the month are tabulated. Also included are the corresponding baseline values, which are the differences between the absolute measurements and the variometer measurements of D , H and Z (in the sense absolute–variometer). These are also plotted (markers) along with the derived preliminary daily baseline values (line) throughout the year. Daily mean differences between the measured absolute F and the F computed from the baseline corrected H and Z values are plotted in the fourth panel (in the sense measured–derived). The bottom panel shows the daily mean temperature in the fluxgate chamber.

4.2 Summary magnetograms

Small-scale magnetograms are plotted which allow the month's data to be viewed at a glance. They are plotted 16 days to a page and show the one-minute variations in D , H and Z . The scales are shown on the right-hand side of the page. On disturbed days the scales are multiplied by a factor, which is indicated above the panel for that day. The variations are centred on the monthly mean value, shown on the left side of the page.

4.3 Magnetograms

The daily magnetograms are plotted using one-minute values of D , H and Z from the fluxgate sensors, with any gaps filled using back-up data. The magnetograms are plotted to a variable scale; scale bars are shown to the right of each plot. The absolute level (the monthly mean value) is indicated on the left side of the plots.

4.4 Hourly Mean Value Plots

Hourly mean values of D , H and Z for the past 12 months are plotted in 27-day segments corresponding to the Bartels solar rotation number. Magnetic disturbances associated with active regions and/or coronal holes on the Sun may recur after 27 days: the same is true for geomagnetically

quiet intervals. Plotting the data in this way highlights this recurrence. Diurnal variations are also clear in these plots and the amplitude changes throughout the year highlight the seasonal changes. Longer term secular variation is also illustrated.

4.5 Daily and Monthly Mean Values

Daily mean values of D , H , Z and F are plotted throughout the year. In addition, a table of monthly mean values of all the geomagnetic elements is provided. These values depend on accurate specification of the fluxgate sensor baselines. It is anticipated that these provisional values will not be altered by more than a few nT or tenths of arcminutes before being made definitive at the end of the year.

5. Conditions of Use

The data presented in this bulletin are provided for personal, academic, educational, non-commercial research or other non-commercial use and are not for sale or distribution to third parties without written permission from BGS.

Reproduction of any part of this bulletin should be accompanied by the statement: 'Reproduced with the permission of the British Geological Survey ©NERC. All rights Reserved'. Publications making use of the data should include an acknowledgment statement of the form: 'The results presented in this paper rely on the data collected at Port Stanley magnetic observatory, operated by the British Geological Survey.'

Commercial users can contact the geomagnetism team for information on the range of applications and services offered. Full contact details are available at www.geomag.bgs.ac.uk/contactus/staff

PORT STANLEY OBSERVATORY

ABSOLUTE OBSERVATIONS

Date	Day Number	Declination			Inclination		Total Field		Horizontal Intensity		Vertical Intensity		Observer
		Time (UT)	Absolute (°)	Baseline (°)	Time (UT)	Absolute (°)	Site difference (nT)	Absolute corrected (nT)	Absolute (nT)	Baseline (nT)	Absolute (nT)	Baseline (nT)	
04-Mar-12	64	13:01	3.2223	3.6683	13:07	-49.5475	-31.4	28590.6	18550.1	18846.5	-21755.9	-21943.0	NB
04-Mar-12	64	13:15	3.2202	3.6667	13:21	-49.5500	-31.4	28590.2	18548.9	18847.8	-21756.3	-21941.9	NB
11-Mar-12	71	13:12	3.2155	3.6650	13:24	-49.5793	-31.4	28583.8	18533.6	18846.1	-21760.9	-21943.2	TS
11-Mar-12	71	13:33	3.2223	3.6683	13:45	-49.5552	-31.4	28582.7	18542.0	18849.2	-21752.4	-21940.3	TS
11-Mar-12	71	14:03	3.2454	3.6667	14:14	-49.5748	-31.4	28582.9	18534.8	18843.9	-21758.9	-21944.8	AS
11-Mar-12	71	14:25	3.2645	3.6683	14:34	-49.5714	-31.4	28581.2	18534.9	18846.8	-21756.4	-21942.4	AS
12-Mar-12	72	12:56	3.2923	3.6750	13:05	-49.5780	-31.4	28618.5	18556.6	18844.2	-21786.9	-21944.4	TS
12-Mar-12	72	13:12	3.2789	3.6750	13:18	-49.5781	-31.4	28612.9	18552.9	18845.8	-21782.7	-21943.2	TS
12-Mar-12	72	13:26	3.2638	3.6733	13:34	-49.5706	-31.4	28608.3	18552.8	18846.1	-21776.8	-21942.6	AS
12-Mar-12	72	13:41	3.2696	3.6717	13:48	-49.5673	-31.4	28607.9	18553.8	18846.8	-21775.4	-21942.1	AS
25-Mar-12	85	15:38	3.2827	3.6717	15:45	-49.6069	-31.4	28581.7	18521.8	18844.3	-21768.3	-21943.6	NB
25-Mar-12	85	15:52	3.2958	3.6683	15:59	-49.6064	-31.4	28581.0	18521.5	18844.9	-21767.6	-21943.0	NB

Falkland Island 2012

Declination : Absolute - Variometer (markers) and Baseline Applied (line)

220.3

5min

Horizontal Intensity : Absolute - Variometer (markers) and Baseline Applied (line)

18847

20nT

Vertical Intensity : Absolute - Variometer (markers) and Baseline Applied (line)

-21943

20nT

Total Intensity : Absolute (Proton F) - Baseline Adjusted Variometer (derived F)

31.4


20nT

Temperature Inside Variometer Hut

23.4

20°C


Jan Feb Mar Apr May Jun Jul Aug Sep Oct Nov Dec




Falkland Islands

March

2012






Falkland Islands March 2012

Date: 01-03-2012


Falkland Islands

Day number: 061



Date: 02-03-2012


Day number: 062



Date: 03-03-2012


Falkland Islands

Day number: 063



Date: 04-03-2012


Day number: 064



Date: 05-03-2012


Falkland Islands

Day number: 065



Date: 06-03-2012


Day number: 066



Date: 07-03-2012


Falkland Islands

Day number: 067



Date: 08-03-2012


Day number: 068



Date: 09-03-2012


Falkland Islands

Day number: 069



Date: 10-03-2012


Day number: 070



Date: 11-03-2012


Falkland Islands

Day number: 071



Date: 12-03-2012


Day number: 072



Date: 13-03-2012


Falkland Islands

Day number: 073



Date: 14-03-2012


Day number: 074



Date: 15-03-2012


Falkland Islands

Day number: 075



Date: 16-03-2012


Day number: 076



Date: 17-03-2012


Falkland Islands

Day number: 077



Date: 18-03-2012


Day number: 078



Date: 19-03-2012


Falkland Islands

Day number: 079



Date: 20-03-2012


Day number: 080



Date: 21-03-2012


Falkland Islands

Day number: 081



Date: 22-03-2012


Day number: 082



Date: 23-03-2012


Falkland Islands

Day number: 083



Date: 24-03-2012


Day number: 084



Date: 25-03-2012


Falkland Islands

Day number: 085



Date: 26-03-2012


Day number: 086



Date: 27-03-2012


Falkland Islands

Day number: 087



Date: 28-03-2012


Day number: 088



Date: 29-03-2012


Falkland Islands

Day number: 089



Date: 30-03-2012


Day number: 090




Date: 31-03-2012

Falkland Islands


Day number: 091




Falkland Islands Observatory: Declination (degrees)




Falkland Islands Observatory: Horizontal Intensity (nT)



Falkland Islands Observatory: Vertical Intensity (nT)



Falklands Is Observatory 2012



Monthly Mean Values for Port Stanley Observatory 2012

Month	<i>D</i>	<i>H</i>	<i>I</i>	<i>X</i>	<i>Y</i>	<i>Z</i>	<i>F</i>
January	3° 19.2′	18587 nT	-49° 29.8′	18556 nT	1077 nT	-21760 nT	28617 nT
February	3° 18.8′	18571 nT	-49° 31.3′	18540 nT	1074 nT	-21760 nT	28607 nT
March	3° 17.9′	18544 nT	-49° 34.0′	18514 nT	1067 nT	-21763 nT	28593 nT

Note

i. The values shown here are provisional.