



British
Geological
Survey



MONTHLY MAGNETIC BULLETIN

Port Stanley Observatory

December 2023



MONTHLY MAGNETIC BULLETIN

Port Stanley Observatory magnetic data

1. Introduction

Port Stanley Observatory was installed by the British Geological Survey (BGS) with financial support from a consortium of oil companies and became operational in February 1994.

This bulletin is published to provide rapid access to the provisional geomagnetic observatory results. The information is freely available for personal, academic, educational and non-commercial research or use. Magnetic observatory data are presented as a series of plots of one-minute, hourly and daily values, followed by tabulations of monthly values. The operation of the observatory and presentation of data are described in the rest of this section.

Enquiries about the data should be addressed to:

Geomagnetism Team
British Geological Survey
The Lyell Centre
Research Avenue South
Edinburgh EH14 4AP
Scotland, UK

Tel: +44 (0) 131 667 1000

Email: enquiries@bgs.ac.uk

Internet: geomag.bgs.ac.uk

2. Position

Port Stanley Observatory, one of the geomagnetic observatories maintained and operated by the British Geological Survey (BGS), is situated on a site at Sapper Hill near Port Stanley in the Falkland Islands. In 2013 it was necessary to establish a new position for the observatory absolute pillar due to degradation in the quality of absolute observations caused by anthropogenic noise. Following an overlap period of at least six months, the observatory results relate to the new position from 1 January 2014:

Old observatory co-ordinates are (Feb 1994 to Dec 2013):

Geographic: $51^{\circ} 42'15"S$ $302^{\circ} 06'24"E$

Height above mean sea level: 135 m

New observatory co-ordinates are (Jan 2014 to present):

Geographic: $51^{\circ} 42'18.0"S$ $302^{\circ} 06'25.2"E$

Geomagnetic: $42^{\circ} 40'26.4"S$ $012^{\circ} 24'25.2"E$

Height above mean sea level: 130 m

The geographical coordinates are measured by a handheld GPS device, which uses WGS84 as the reference coordinate system. The height above MSL is determined from the best available contour maps. The geomagnetic co-ordinates are approximations, calculated using the 13th generation International Geomagnetic Reference Field (IGRF) at epoch 2022.5. On-line access to models (including IGRF), charts and navigational data are available at geomag.bgs.ac.uk/data_service/models_compass/home

3. The observatory operation

3.1 GDAS

The observatory operates under the control of the Geomagnetic Data Acquisition System (GDAS), which was developed by BGS staff, installed and became operational in August 2002. The data acquisition software, running on QNX operated computers, controls the data logging and the communications.

There are two sets of sensors used for making magnetic measurements. A tri-axial linear-core fluxgate magnetometer, manufactured by the Danish Meteorological Institute, is used to measure the variations in the horizontal (H) and vertical (Z) components of the field. The third sensor is oriented perpendicular to these, and measures variations, which are proportional to the changes in declination (D). Measurements are made at a rate of 1 Hz.

In addition to the fluxgate sensors there is a proton precession magnetometer (PPM) making measurements of the absolute total field intensity (F) at a rate of 0.1Hz.

The raw unfiltered data are retrieved automatically via Internet connections to the BGS office in Edinburgh in near real-time. The fluxgate data are filtered to produce one-minute values using a 61-point cosine filter and the total field intensity samples are filtered using a 7-point cosine filter. The one-minute values provide input for various data products, available on-line at geomag.bgs.ac.uk/data_service/home

3.2 Absolute observations

The GDAS fluxgate magnetometers accurately measure variations in the components of the geomagnetic field, but not the absolute magnitudes. Two sets of absolute measurements of the field are made manually twice per month. A fluxgate sensor mounted on a theodolite is used to determine D and inclination (I); the GDAS PPM measurements, with a site difference correction applied, are used for F . The absolute observations are used in conjunction with the GDAS variometer measurements to produce a continuous record of the absolute values of the geomagnetic field elements as if they had been measured at the observatory reference pillar.

4. Observatory results

The data presented in the bulletin are in the form of plots and tabulations described in the following sections.

4.1 Absolute observations

The absolute observation measurements made during the month are tabulated. Also included are the corresponding baseline values, which are the differences between the absolute measurements and the variometer measurements of D , H and Z (in the sense absolute-variometer). These are also plotted (markers) along with the derived preliminary daily baseline values (line) throughout the year. Daily mean differences between the measured absolute F and the F computed from the baseline corrected H and Z values are

plotted in the fourth panel (in the sense measured-derived). The bottom panel shows the daily mean temperature in the fluxgate chamber.

4.2 Summary magnetograms

Small-scale magnetograms are plotted which allow the month's data to be viewed at a glance. They are plotted 16 days to a page and show the one-minute variations in D , H and Z . The scales are shown on the right-hand side of the page. On disturbed days the scales are multiplied by a factor, which is indicated above the panel for that day. The variations are centred on the monthly mean value, shown on the left side of the page.

4.3 Magnetograms

The daily magnetograms are plotted using one-minute values of D , H and Z from the fluxgate sensors, with any gaps filled using back-up data. The magnetograms are plotted to a variable scale; scale bars are shown to the right of each plot. The absolute level (the monthly mean value) is indicated on the left side of the plots.

4.4 Hourly mean value plots

Hourly mean values of D , H and Z for the past 12 months are plotted in 27-day segments corresponding to the Bartels solar rotation number. Magnetic disturbances associated with active regions and/or coronal holes on the Sun may recur after 27 days: the same is true for geomagnetically quiet intervals. Plotting the data in this way highlights this recurrence. Diurnal variations are also clear in these plots and the amplitude changes throughout the year highlight the seasonal changes. Longer term secular variation is also illustrated.

4.5 Daily and monthly mean values

Daily mean values of D , H , Z and F are plotted throughout the year. In addition, a table of monthly mean values of all the geomagnetic elements is provided. These values depend on accurate specification of the fluxgate sensor baselines. It is anticipated that these provisional values will not be altered by more than a few nT or tenths of arcminutes before being made definitive at the end of the year.

5. Conditions of use

The data presented in this bulletin are provided for personal, academic, educational, non-commercial research or other non-commercial use and are not for sale or distribution to third parties without written permission from BGS.


Reproduction of any part of this bulletin should be accompanied by the statement: 'Reproduced with the permission of the British Geological Survey ©UKRI. All rights Reserved'. Publications making use of the data should include an acknowledgment statement of the form: 'The results presented in this paper rely on the data collected at Port Stanley magnetic observatory, operated by the British Geological Survey.'

Commercial users can contact the geomagnetism team for information on the range of applications and services offered. Full contact details are available at geomag.bgs.ac.uk/contactus/staff

Port Stanley Observatory

Absolute observations


Port Stanley 2023




Falkland Islands

December

2023






Date: 01-12-2023


Falkland Islands

Day number: 335



Date: 02-12-2023


Day number: 336



Date: 03-12-2023


Falkland Islands

Day number: 337



Date: 04-12-2023


Day number: 338



Date: 05-12-2023


Falkland Islands

Day number: 339



Date: 06-12-2023


Day number: 340



Date: 07-12-2023


Falkland Islands

Day number: 341



Date: 08-12-2023


Day number: 342



Date: 09-12-2023


Falkland Islands

Day number: 343



Date: 10-12-2023


Day number: 344



Date: 11-12-2023


Falkland Islands

Day number: 345



Date: 12-12-2023


Day number: 346



Date: 13-12-2023


Falkland Islands

Day number: 347



Date: 14-12-2023


Day number: 348



Date: 15-12-2023


Falkland Islands

Day number: 349



Date: 16-12-2023


Day number: 350



Date: 17-12-2023


Falkland Islands

Day number: 351



Date: 18-12-2023


Day number: 352



Date: 19-12-2023


Falkland Islands

Day number: 353



Date: 20-12-2023


Day number: 354



Date: 21-12-2023


Falkland Islands

Day number: 355



Date: 22-12-2023


Day number: 356



Date: 23-12-2023


Falkland Islands

Day number: 357



Date: 24-12-2023


Day number: 358



Date: 25-12-2023


Falkland Islands

Day number: 359



Date: 26-12-2023


Day number: 360



Date: 27-12-2023


Falkland Islands

Day number: 361



Date: 28-12-2023


Day number: 362



Date: 29-12-2023


Falkland Islands

Day number: 363



Date: 30-12-2023


Day number: 364




Date: 31-12-2023

Falkland Islands


Day number: 365




Falkland Islands Observatory: Declination (degrees)



Falkland Islands Observatory: Horizontal Intensity (nT)




Falkland Islands Observatory: Vertical Intensity (nT)



Falklands Is Observatory

2023



Monthly mean values for Port Stanley Observatory 2023

Month	D	H	I	X	Y	Z	F
January	2° 25.3'	17821 nT	-50° 24.6'	17805 nT	753 nT	-21550 nT	27963 nT
February	2° 25.0'	17801 nT	-50° 26.5'	17785 nT	751 nT	-21550 nT	27951 nT
March	2° 24.2'	17784 nT	-50° 28.5'	17768 nT	746 nT	-21554 nT	27944 nT
April	2° 23.6'	17780 nT	-50° 28.7'	17764 nT	742 nT	-21552 nT	27940 nT
May	2° 23.4'	17776 nT	-50° 29.2'	17760 nT	741 nT	-21554 nT	27938 nT
June	2° 22.9'	17777 nT	-50° 28.7'	17762 nT	739 nT	-21549 nT	27935 nT
July	2° 22.8'	17775 nT	-50° 28.7'	17760 nT	738 nT	-21546 nT	27932 nT
August	2° 22.5'	17771 nT	-50° 28.9'	17756 nT	736 nT	-21544 nT	27928 nT
September	2° 22.4'	17759 nT	-50° 29.8'	17744 nT	735 nT	-21542 nT	27918 nT
October	2° 21.6'	17763 nT	-50° 29.0'	17748 nT	732 nT	-21535 nT	27916 nT
November	2° 21.8'	17758 nT	-50° 29.7'	17743 nT	733 nT	-21538 nT	27915 nT
December	2° 21.3'	17762 nT	-50° 29.1'	17747 nT	730 nT	-21536 nT	27916 nT

Note:

- i. The values shown here are provisional.